

Master Medicine Protocol for the Administration of Spikevax® (COVID-19 Vaccine Moderna) to Vaccine Recipients

This medicine protocol is a specific written instruction for the administration of Spikevax® (COVID-19 Vaccine Moderna) to vaccine recipients by healthcare professionals included in Statutory Instruments S.I. No. 698 of 2020, S.I. No. 81 of 2021 and S.I. No. 245 of 2021 who are registered with their respective regulatory body and students in healthcare professions included in S.I. No. 245 of 2021. This medicine protocol is valid for the 2020/2021 HSE COVID-19 Vaccination Programme. This medicine protocol enables the healthcare professionals and students described above who are employed in the voluntary and statutory services of the Health Service Executive (HSE) who have undertaken the required education and training programmes to administer Spikevax® (COVID-19 Vaccine Moderna) to vaccine recipients, with reference to guidelines and guidance from National Immunisation Advisory Committee (NIAC), National Immunisation Office (NIO), HSE and in accordance with the Summary of Product Characteristics for Spikevax® (COVID-19 Vaccine Moderna) as detailed by the European Medicines Agency (EMA).

- National Immunisation Advisory Committee *Immunisation Guidelines for Ireland* Dublin: Royal College of Physicians Ireland (Online Update available at <http://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/>)
- National Immunisation Office (2020) *Clinical Guidance for COVID-19 Vaccinations* (available at <https://www.hse.ie/eng/health/immunisation/hcpinfo/covid19vaccineinfo4hps/clinicalguidance.pdf>)
- Summary of Product Characteristics <https://www.ema.europa.eu/en/medicines/human/EPAR/covid-19-vaccine-moderna#product-information-section>

A medicine protocol has been defined as follows: written directions that allow for the supply and administration of a named medicinal product by specified healthcare professionals and students in identified clinical situations. A medicine protocol involves the authorisation of the healthcare professional or student to supply and administer a medicine to groups of patients in a defined situation meeting specific criteria and who may not be individually identified before presentation for treatment. An individually named prescription is not required for the supply and administration of medication when a medication protocol is in effect

The HSE has developed this medicines protocol to facilitate the delivery of COVID-19 immunisation in line with NIAC recommendations, Department of Health (DoH) and HSE policy.

The professional groups and students using this protocol must ensure that the protocol is organisationally authorised by an appropriate authorising person, relating to the professional or student cohort of vaccinators by whom the vaccine is to be administered, including requirements of registration, training and assessment of competency.

Medicine Protocol for the Administration of Spikevax® (COVID-19 Vaccine Moderna) to vaccine recipients

Document reference number:	NIO 001.2
1.0 Critical Elements	
Name of Organisation where medicine protocol applies	Health Service Providers across the voluntary and statutory services of the Health Service Executive (HSE), non-HSE healthcare facilities and mass vaccination clinic venues. This Medicine Protocol applies to: Registered healthcare professionals included in S.I. 698 of 2020, S.I. 81 of the 2021, S. I. No. 245 and employed in the voluntary and statutory services of the Health Service Executive (HSE) and students in healthcare professions included in S.I. No. 245 of 2021 who have undertaken the required education and training programmes
Date the medicine protocol comes into effect	January 2021
Date for review of medicine protocol	January 2022
Document prepared by:	The National Immunisation Office (NIO)
Names and Signatures of the employing authority who is authorising the implementation of the medicine protocol <i>“On behalf of the authority employing professionals authorised to administer under this medicine protocol, I have read this medicine protocol and authorise its implementation”</i>	<p>Name: Dr Kevin Kelleher, Assistant National Director, National Office for Public Health/Child Health Strategic Planning and Transformation, HSE</p> <p>Signature: </p> <p>Name: Dr Colm Henry, Chief Clinical Officer, HSE</p> <p>Signature: </p>

2.0 Clinical Criteria	
Clinical Condition for use of the medicine protocol	The clinical condition for which this medicine protocol has been developed is for the immunisation of vaccine recipients (see Inclusion Criteria) against COVID-19.
Circumstances in which the medicine protocol applies	Targeted immunisation programme for vaccine recipients against COVID-19 as identified in the DoH policy based on the NIAC recommendations. The World Health Organisation declared COVID-19 outbreak as a pandemic on 11th March 2020 which is still ongoing.
Inclusion criteria for vaccine recipient using the medicine protocol	<p>Note: Vaccine Recipients who have received Spikevax® (COVID-19 Vaccine Moderna) as a first dose MUST be advised that the second dose is ALSO Spikevax® (COVID-19 Vaccine Moderna) ONLY.</p> <p>Inclusion Criteria:</p> <ul style="list-style-type: none"> ● Active immunisation to prevent COVID-19 caused by SARS-CoV-2 virus, in individuals 18 years of age and older. <p>Precautions:</p> <ul style="list-style-type: none"> ● Acute severe febrile illness defer until recovery ● Those with the following history should receive a viral vector vaccine: <ul style="list-style-type: none"> ● Anaphylaxis after multiple, different drug classes, with no identified allergen (may indicate PEG allergy) ● Anaphylaxis after a vaccine, or a medicine which contained PEG ● Idiopathic anaphylaxis (may indicate PEG allergy) <p>If vaccination is advised for a person with prior anaphylaxis to an unrelated allergen observe for 30 minutes after vaccination.</p> <ul style="list-style-type: none"> ● Vaccination should be deferred until clinical recovery from COVID-19 at least four weeks after diagnosis or onset of symptoms, or four weeks from the ● first PCR positive specimen in those who are asymptomatic ● Vaccination is not contraindicated for those with persisting symptoms post COVID-19 unless there is evidence of recent clinical deterioration ● Individuals with a bleeding disorder or receiving anticoagulant therapy may develop haematomas in intramuscular (IM) injection sites. Prior to vaccination, inform the recipient about this risk. For those with thrombocytopenia (platelet count <50 x 10³/ml) consult the supervising consultant ● Those with inherited coagulopathies who require factor replacement therapy should receive it on the day of vaccination, prior to the IM vaccination. If there is uncertainty about the need for cover, contact the patient's Comprehensive Care Centre ● COVID-19 vaccines and other vaccines may be administered at the same time or at any interval. As it is not known if COVID-19 vaccine reactogenicity is increased with coadministration, vaccines should preferably be given in different limbs. ● Patients with planned immunosuppressive therapy should ideally complete vaccination two weeks before treatment. The recommended minimum interval may be used. <p>Pregnancy:</p> <ul style="list-style-type: none"> ● Women who are at less than 14 weeks or more than 36 weeks of gestation should not receive the vaccine

	<ul style="list-style-type: none"> Pregnant women who are between 14 weeks and 36 weeks of gestation and wish to receive the vaccine should confirm they have consulted with their obstetric care giver (Obstetrician or GP) and decided to receive the vaccine. <p>Breastfeeding:</p> <ul style="list-style-type: none"> There is no known reason for vaccine recipients to avoid breastfeeding. Breastfeeding mothers should be vaccinated according to their risk grouping.
Exclusion criteria for vaccine recipient using the medicine protocol	<p>Spikevax® (COVID-19 Vaccine Moderna) should not be given under this medicine protocol if the vaccine recipient has:</p> <ul style="list-style-type: none"> Anaphylaxis (serious systemic allergic reaction requiring medical intervention) following a previous dose of the vaccine or any of its constituents (including polyethylene glycol (PEG)). Anaphylaxis following another mRNA vaccine. Those with a contraindication to one mRNA COVID-19 vaccine should not receive another authorised mRNA vaccine.
Actions to be taken for those who are excluded from the medicine protocol	<ul style="list-style-type: none"> Refer to/discuss with the relevant Medical Practitioner/clinical lead/lead vaccinator for an individual medical assessment. Those with a contraindication to one mRNA COVID-19 vaccine should not receive another authorised mRNA vaccine. Consideration may be given to viral vector vaccination which should be given after an interval of at least 28 days. The Medical Practitioner/clinical lead/lead vaccinator can consider referring the individual to an allergist/Immunologist for a further assessment. Document action in clinical record or IT system Where Spikevax® (COVID-19 Vaccine Moderna) is prescribed following medical assessment, the vaccinator may administer the vaccine within his/her scope of practice. <p>Note: In determining their scope of practice, vaccinators must make judgements about their competency to carry out a role or activity in accordance with the guidance from their regulator</p>
Action to be followed for vaccine recipients who do not wish to receive the vaccine	<p>Advise of the risks of not having the vaccine, including risk of possible severe COVID-19 disease.</p> <p>Advise regarding minimization of risk.</p>
Description of circumstances and referral arrangements when further advice or consultation is required	<p>Refer to/discuss with relevant Medical Practitioner /clinical lead/ lead vaccinator if the vaccine recipient had previous adverse reaction or other clinical concerns as outlined in Exclusion Criteria.</p>

<p>Documentation required to support implementation of the medicine protocol</p>	<ul style="list-style-type: none"> • Check for and ensure consent has been obtained • Vaccine Information Leaflets • Patient held record cards if available • Health Products Regulatory Authority Adverse Reaction Reporting forms • National Incident Management System Form NIRF-01-v11 available at: https://www.hse.ie/eng/about/qavd/incident-management/nirf-01-v11-person.pdf <p>It is the responsibility of each vaccinator to be familiar with the appropriate documentation to support the safe administration of Spikevax® (COVID-19 Vaccine Moderna) which includes the following:</p> <ul style="list-style-type: none"> • Medicine Protocol for the Administration of Spikevax® (COVID-19 Vaccine Moderna) to vaccine recipients • Treatment of anaphylaxis in the community. National Immunisation Advisory Committee, Immunisation Guidelines for Ireland. https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/anaphylaxis.pdf https://www.hse.ie/eng/health/immunisation/hcpinfo/fluinfo/adrenalineprotocol.pdf • Clinical Guidance for Covid-19 Vaccination, available at https://www.hse.ie/eng/health/immunisation/hcpinfo/covid19vaccineinfo4hours/clinicalguidance.pdf <p>COVID-19 chapter from NIAC immunisation Guidelines for Ireland (2020) available at https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/covid19.pdf</p>
---	---

<p>3.0 Name of Medicine</p>	<p>Spikevax® (COVID-19 Vaccine Moderna)</p>
<p>Dose & Route of administration</p>	<ul style="list-style-type: none"> • The dose is 0.5ml, 2 doses 28 days apart • Route of administration: Intramuscular (IM) • Site: The preferred site is the deltoid muscle • If the interval between doses is longer than 28 days, the second dose should be given as soon as possible. The course does not need to be restarted • If the second dose was given between 24 and 27 days after the first dose, it is a valid dose • If the second dose is given before 24 days, this is not considered a valid vaccine. A third dose should be given 28 days after the second (invalid) vaccine. • Do not inject the vaccine intravascularly, subcutaneously or intradermally
<p>Link to Medicine Details of product information and other data including instructions for supply and administration is available from the European Medicines Agency (EMA)</p>	<p>Link to Summary of Product Characteristics and Patient Information Leaflet available at: https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-moderna-product-information_en.pdf</p>

<p>Potential adverse reactions and procedures for treatment of same</p>	<p>Following administration of the vaccine, the vaccine recipient should be advised to remain seated in the post vaccination observation area to enable monitoring of any immediate reaction including suspected anaphylactic reaction</p> <ul style="list-style-type: none"> ● Those with no history of anaphylaxis from any cause: 15 minutes ● Those with a history of anaphylaxis from any cause: 30 minutes ● Those with immediate itching, swelling or urticarial reaction at the vaccination site: 30 minutes or longer as clinically indicated <p>The vaccine recipient should be advised to contact relevant medical personnel in the event of adverse reaction occurring following administration of the COVID-19 Vaccine Moderna after the above period of observation.</p>
<p>Procedure for reporting Adverse Drug Reactions to the Health Products Regulatory Authority (HPRA)</p>	<p>The vaccinator should report to the HPRA any suspected adverse reactions, in accordance with criteria outlined by the HPRA. This reporting may be carried out on line at http://www.hpra.ie or through use of the yellow card system which is available in a downloadable format from the HPRA website, or on request from the HPRA.</p> <p>The vaccine recipient’s General Practitioner should be informed of any clinically significant reported adverse reaction.</p> <p>The incident and all actions taken must be promptly recorded in accordance with the <i>Management of a Patient with Anaphylaxis: Treatment in the Community</i> (National Immunisation Advisory Committee 2019), available online at https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/anaphylaxis.pdf</p>
<p>Procedure for the reporting and documentation of errors and near misses involving the medicine</p>	<p>In the case of medication errors that directly involve the vaccine recipient, i.e. wrong medication/dose/route being administered or another medication error, the vaccinator must remain with the person and closely monitor them for any adverse reactions.</p> <p>The recipient should be reviewed by the relevant medical practitioner/clinical lead/ lead vaccinators and the Vital signs to be recorded.</p> <p>The incident must be reported to the relevant line manager/person in charge as soon as possible.</p> <p>The incident and all actions taken must be recorded and the relevant National Incident Management Report Form (NIRF) completed as soon as is practicable after the event occurs and within one working day. (National Incident Report Form (NIRF 01 – V11)) (2020) available at: https://www.hse.ie/eng/about/qavd/incident-management/nirf-01-v11-person.pdf The vaccine recipient and/or significant others should be informed of the incident.</p> <p>An incident report form must be completed by the vaccinators and forwarded to local or regional Risk Manager as per local policy.</p> <p>Any suspected adverse reactions associated with medication errors should be reported to the HPRA as outlined above.</p>

Resources and equipment required	<ul style="list-style-type: none"> ● A multidose vial of Spikevax® (COVID-19 Vaccine Moderna) ● 1 ml/2ml/2.5ml syringe, 23/25 gauge needle for IM administration ● Fridge/Cooler box with data logger with external temperature monitoring display to maintain cold chain temperature between +2° to +8°C ● Disposable kidney dishes/trays ● 70% alcohol swabs (for sterilizing vials) ● Gauze swabs, tape/plasters ● Sharps bins, and bins for the disposal of healthcare risk and non-risk waste ● Alcohol hand sanitiser ● Access to telephone ● Resuscitation equipment and drugs in accordance with Anaphylaxis: Treatment in the Community (National Immunisation Advisory Committee, 2019) available at https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/anaphylaxis.pdf ● Safe storage areas for medicines and equipment ● Current COVID-19 Vaccine Moderna medicine protocol
Audit process to identify appropriate use of the medicine protocol or unexpected outcomes	All documentation will be held for review and audit purposes as per local/national agreement.
4.0 Information for vaccine recipient	
Advice to be given to the vaccine recipient before treatment	<p>Vaccine Information material must be supplied to the vaccine recipient prior to administration of the vaccine.</p> <p>Before Treatment Check and confirm that consent has been obtained</p> <p>Discuss the Spikevax® (COVID-19 Vaccine Moderna) and the importance of protecting their health. Inform vaccine recipient that patient information leaflet is available online at https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-moderna-product-information_en.pdf</p> <p>Discuss potential side effects as below Side effects may occur with following frequencies:</p> <p>Local: Very common: injection site pain and swelling Common: injection site erythema, rash and urticaria Uncommon: injection site pruritus.</p> <p>General: Very common: arthralgia, axillary lymphadenopathy (on the side of injection), chills, fatigue, fever, headache, myalgia, nausea, vomiting Rare: acute peripheral facial paralysis, facial swelling (in those with dermatological fillers)</p> <p>A full list of adverse reactions may be found in the Summary of Product Characteristics (SmPC), available at https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-moderna-product-information_en.pdf</p> <p>Individuals may not be protected until at least 14 days after their second dose of the</p>

	vaccine.
--	----------

<p>Advice to be given to the recipient after treatment</p>	<p>After Treatment Discuss potential side effects Following administration of the vaccine, the vaccine recipient should be advised to remain seated in the post vaccination observation area to enable monitoring of any immediate reaction including suspected anaphylactic reaction. Events of anaphylaxis have been reported therefore NIAC recommends the following monitoring for the post-vaccination period:</p> <ul style="list-style-type: none"> ● Close observation for at least 15 minutes is recommended following vaccination ● The second dose of the vaccine should not be given to those who have experienced anaphylaxis to the first dose of COVID-19 Vaccine Moderna. <p>The vaccine recipient should not leave the healthcare facility if they are feeling unwell and must report any side effects to a member of the vaccination team. If required, symptomatic treatment with analgesic and/or anti-pyretic medicinal products (e.g. paracetamol or ibuprofen-containing products) may be used. Ibuprofen is not recommended in pregnancy. If more serious adverse or persistent effects occur, vaccine recipient should be advised to contact their GP/out of hours service.</p>
<p>Details of any necessary follow-up, action and referral arrangements</p>	<p>In the event of an adverse reaction the vaccination team must ensure that all procedures are adhered to as outlined in Section 3.</p>

References

Health Service Executive (2010) *Healthcare Risk Waste Management Segregation Packaging and Storage Guidelines for healthcare Risk Waste*. Dublin: Health Service Executive.

National Immunisation Advisory Committee (2019) *Anaphylaxis: Treatment in the Community*. Available at <https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/anaphylaxis.pdf>

National Immunisation Advisory Committee *Immunisation Guidelines for Ireland (2020)* Dublin: Royal College of Physicians Ireland. Online update available at <http://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/>

National Immunisation Office (2020) *Clinical Guidance for COVID-19 Vaccinations* (available at <https://www.hse.ie/eng/health/immunisation/hcpinfo/covid19vaccineinfo4hps/>)

S.I. No. 81/2021 - Medicinal Products (Prescription and Control of Supply) (Amendment) (No.4) Regulations 2021. Available at <http://www.irishstatutebook.ie/eli/2021/si/81/made/en/pdf>

S.I. No. 698/2020 - Medicinal Products (Prescription and Control of Supply) (Amendment) (No. 7) Regulations 2020. Available at <http://www.irishstatutebook.ie/eli/2020/si/698/made/en/pdf>

Section B Information Specific to Registered Nurses and Registered Midwives for the administration of the COVID-19 vaccines



Seirbhís Sláinte
Níos Fearr
á Forbairt

Building a
Better Health
Service



Office of the
Nursing & Midwifery
Services Director

Statement of Support from Dr Geraldine Shaw, Nursing and Midwifery Services Director, Office of the Nursing and Midwifery Services, HSE

I am delighted to support Registered Nurses and Registered Midwives to administer COVID-19 vaccines under medicine protocol.

Nurses and midwives have a long tradition of supporting vaccination programmes, for example Schools Immunisation Programme, Seasonal Influenza Peer Vaccination Programme and Primary Childhood Immunisation Programme.

The national COVID-19 vaccination programme commenced in December 2020. Statutory Instruments No. 698 of 2020, No. 8 of 2021 and No. 43 of 2021 identify nurses and midwives as professions that can administer named COVID-19 vaccines, subject to approval of an education programme by the regulatory body concerned.

In order to administer the vaccines, registered nurses and registered midwives must be familiar with the most up to date version of the medicine protocols including the content of this section and have completed the *COVID-19 Vaccination Programme for Nurses and Midwives* on HSELand. Nurses and midwives must also have completed the Competency Assessment Form, also included in this section.

I would like to acknowledge the contribution of the nursing and midwifery professions to this very important national initiative.

A handwritten signature in black ink, appearing to read 'G. Shaw'.

Signature

30th March 2021

Date

Professional Qualifications, Training, Experience and Competence Required

<p>Professional qualifications, training, experience and competence required prior to using this medicine protocol / Professional Qualifications :</p> <p>Training, Experience, Competence:</p>	<p>Registered nurse or registered midwife, maintained on the active register maintained by The Nursing and Midwifery Board of Ireland.</p> <p>HSELand education programme titled <i>COVID-19 Vaccination Programme for Nurses and Midwives</i></p> <p>Basic Life Support for Health Care Providers within the last two years.</p> <p>Initial anaphylaxis programme (<i>“National Anaphylaxis Education Programme for Health Care Professionals”</i>) via HSELand followed by a one and a half hour classroom based skills workshop (replacing the previous four hour classroom based programme). Subsequent updates every two years via HSELand Anaphylaxis e-learning programme available at www.hse.ie.</p> <p>The nurse/midwife must complete the <i>Competency Assessment Form</i> to administer the COVID-19 Vaccines.</p> <p>COVAX IBM/Salesforce online programme https://www.hse.ie/eng/health/immunisation/hcpinfo/hsecovid19vms.html</p> <p>Recommended:</p> <p><i>Storing and Managing Vaccines</i> www.hseland.ie</p>
--	--

Supporting Documents for Registered Nurses and Registered Midwives

An Bord Altranais (2007) *Guidance to Nurses and Midwives on Medication Management* Dublin: An Bord Altranais
Health Service Executive (2019) *Directions for Nurses and Midwives for the Management of a Patient who Develops Anaphylaxis or Suspected Anaphylaxis incorporating Medicine Protocol for the Administration of Epinephrine (Adrenaline) Injection BP 1:1,000 by intramuscular injection by nurses and midwives for the management of a patient with anaphylaxis or suspected anaphylaxis*. Dublin: Health Service Executive

Government of Ireland (2020) *Statutory Instruments Number 698 of 2020*. Dublin: Stationery Office

Government of Ireland (2021) *Statutory Instruments Number 8 of 2021*. Dublin: Stationery Office

Government of Ireland (2021) *Statutory Instruments Number 43 of 2021*. Dublin: Stationery Office

Health Service Executive (2010) *Healthcare Risk Waste Management Segregation Packaging and Storage Guidelines for healthcare Risk Waste*. Dublin: Health Service Executive.

Nursing and Midwifery Board of Ireland (2014) *Code of Professional Conduct and Ethics for Registered Nurses and Registered Midwives*. Dublin: Nursing and Midwifery Board of Ireland available at:

<http://www.nmbi.ie/Standards-Guidance/Code>.

Nursing and Midwifery Board of Ireland (2015) *Practice Standards for Midwives*. Dublin: Nursing and Midwifery Board of Ireland available at: <http://www.nmbi.ie/Standards-Guidance/Midwives-Standards>.

Nursing and Midwifery Board of Ireland (2015) *Recording Clinical Practice. Guidance to Nurses and Midwives*. Dublin: Nursing and Midwifery Board of Ireland available at: <http://www.nmbi.ie/Standards-Guidance/More-Standards-Guidance/Recording-Clinical-Practice>

Nursing and Midwifery Board of Ireland (2015) *Scope of Nursing and Midwifery Practice Framework*. Dublin: Nursing and Midwifery Board of Ireland available at: <http://www.nmbi.ie/Standards-Guidance/Scope-of-Practice/Nursing-Practise-Scope-Definition>

Nursing and Midwifery Board of Ireland (2020) *Guidance for Registered Nurses and Midwives on Medication Administration*. Dublin: Nursing and Midwifery Board of Ireland, available at: <http://www.nmbi.ie>

Competency Assessment Form



Seirbhís Sláinte
Níos Fearr
á Forbairt

Building a
Better Health
Service



Office of the
Nursing & Midwifery
Services Director

NAME:

Self-Assessment of Competency to Administer COVID-19 Vaccine under Medicine Protocol

Domain of Practice	Critical Element	Competent Date/ Initials	Needs Practice Date/ Initials	Needs Theory Date/ Initials
1	I understand the role and function of medicine protocols in the context of NMBI guidelines in relation to: <ul style="list-style-type: none"> The Code of Professional & Ethical Conduct Scope of Nursing and Midwifery Practice Guidance to Nurses and Midwives on Medication Management NIAC Immunisation Guidelines for Ireland. 			
2	I practice within my scope of practice to undertake administration of COVID-19 Vaccines under medicine protocol.			
3	I have undertaken the <i>COVID-19 Vaccination Programme for Nurses and Midwives</i> on HSE LanD.			
4	I have attended Basic Life Support for Health Care Providers within the last two years.			
5	I am competent in safe injection technique.			
6	I have attended an approved Anaphylaxis education programme and I am familiar with the current medicine protocol on the administration of Epinephrine by RNs/RMs.			
7	I can outline the inclusion/ exclusion criteria for administering COVID-19 Vaccine under the named medicine protocol.			
8	I can refer to/discuss those that are meeting the exclusion criteria to the relevant medical practitioner for an individual medical assessment as per medicine protocol.			
9	I am familiar with the documentation required to support implementation of the medicine protocol to ensure safe administration of COVID-19 Vaccine.			
10	In assessing suitability for vaccination I can undertake a clinical assessment of individuals within the scope of the medicine protocol.			
11	I can provide information regarding COVID-19 Vaccine, benefits and side effects to vaccine recipients.			
12	I am aware of the procedure for treatment and reporting of potential adverse reactions.			
13	I understand the procedure for reporting and documentation of medicine errors/ near misses.			
14	I dispose of all equipment and sharps in accordance with guidance for Healthcare Risk Waste (HSE, 2010).			
15	I am aware of and comply with the guidance on vaccine storage and handling including the maintenance of the cold chain in accordance with national and local policies.			

16	<p>I have undertaken the following HSEland/online programmes:</p> <ul style="list-style-type: none"> ● AMRIC Aseptic Technique www.hseland.ie ● AMRIC Hand Hygiene www.hseland.ie ● GDPR guidelines www.hseland.ie ● COVAX IBM/Salesforce online programme https://www.hse.ie/eng/health/immunisation/hcpinfo/hsecovid19vms.html 			
----	--	--	--	--

I have sufficient theoretical knowledge and practice to undertake vaccination under this medicine protocol independently, and I acknowledge my responsibility to maintain my own competence in line with the Scope of Nursing and Midwifery Practice and current best evidence.

Registered Nurse/Midwife Signature: _____ Date: _____

If any deficits in theory and/or clinical practice are identified, the nurse/midwife must discuss with relevant Line Manager and implement appropriate action plan to achieve competency within an agreed time frame.

<p>Action Plan (for use if needed to reach competencies outlined) Action necessary to achieve competency:</p> <p>.....</p> <p>.....</p> <p>.....</p> <p>.. Date to be achieved:.....</p> <p>Supporting evidence of measures taken to achieve competency:</p> <p>.....</p> <p>.....</p> <p>Nurse/Midwife signature:</p> <p style="text-align: right;">ate: _____</p> <p>Line Manager signature</p> <p style="text-align: right;">ate: _____</p>	
---	--

Nursing and Midwifery Board of Ireland Statement of Support 2021

