

## Tamoxifen Monotherapy

### INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Adjuvant treatment of oestrogen receptor positive breast cancer in pre- or post-menopausal women.	C50	00253a	CDS
Treatment of oestrogen receptor positive advanced breast cancer in pre- or post-menopausal women	C50	00253b	CDS

### TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

Tamoxifen is administered orally once daily continuously during treatment.

**Duration of treatment will be determined by the prescribing Consultant and depends on disease progression or unacceptable toxicity.**

Drug	Dose	Route	Diluent & Rate	Cycle
Tamoxifen	20 mg daily	PO	NA	Continuous for specified duration or until disease progression or unacceptable toxicity
Tablet should be swallowed whole. Can be taken with food or on an empty stomach with a glass of water If nausea develops, tamoxifen may be taken with or after food or at night. If patient vomits within a few hours of taking the drug, do <b>not</b> repeat the dose				
Missed doses should not be replaced, normal dosing should be resumed at the next scheduled daily dose				

### ELIGIBILITY:

- Indications as above

### EXCLUSIONS:

- Hypersensitivity to tamoxifen or any of the excipients.
- Hormone receptor-negative.
- Pregnancy
- Patients with a history of significant thromboembolic disease

### PRESCRIPTIVE AUTHORITY:

Medical oncologist or General Practitioner under direction of plan written by medical oncologist.

### TESTS:

#### Baseline tests:

- FBC, renal and liver profile

#### Regular tests:

- None unless clinically indicated

NCCP Regimen: Tamoxifen Monotherapy	Published: 01/11/2014 Review: 10/11/2025	Version number: 4a
Tumour Group: Breast NCCP Regimen Code: 00253	IHS/ISMO Contributor: Prof Maccon Keane	Page 1 of 3
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## Disease monitoring:

- Metastatic disease: Disease monitoring/assessment should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.
- Adjuvant treatment: No routine tests required

## DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant
- Intolerant or serious complications during tamoxifen therapy. (Note: Post-menopausal patients may be switched to aromatase inhibitor therapy for a total of 5 years of adjuvant hormonal therapy).

## SUPPORTIVE CARE:

**EMETOGENIC POTENTIAL:** Minimal Risk (Refer to local policy).

**PREMEDICATIONS:** Not usually required

## OTHER SUPPORTIVE CARE:

Not usually required

## ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS

*The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.*

- **Myelosuppression:** Mild myelosuppression with transient thrombocytopenia may occur rarely. The association with tamoxifen is uncertain.
- **Endometrial Cancer:** Annual gynecological examinations are recommended. Pelvic complaints, such as unusual vaginal bleeding, require prompt evaluation.
- **Ocular Toxicity:** Ocular toxicity is rare and may occur after only a few weeks of therapy, although it is more common with prolonged treatment. Ophthalmologic examination is recommended if visual disturbances occur.
- **Thromboembolism:** Tamoxifen is associated with an increased risk of thromboembolism that is comparable to estrogen replacement therapy.
- **Hepatotoxicity:** While hepatotoxicity is rare and usually presents as elevated hepatic enzymes, more serious liver abnormalities have been reported.
- **Ovulation Induction:** Tamoxifen may induce ovulation in pre- and peri-menopausal women. Barrier forms of contraception are recommended.
- **Hyperlipidemia:** Elevations in cholesterol and triglycerides may occur in patients with pre-existing hyperlipidemias.

## DRUG INTERACTIONS:

- Current drug interaction databases should be consulted for more information.
  - Medications that inhibit CYP2D6 (e.g., paroxetine, fluoxetine, quinidine, bupropion) should be avoided as they may decrease the efficacy of tamoxifen.
- Co-administration of tamoxifen and aromatase inhibitors should be avoided.

NCCP Regimen: Tamoxifen Monotherapy	Published: 01/11/2014 Review: 10/11/2025	Version number: 4a
Tumour Group: Breast NCCP Regimen Code: 00253	IHS/ISMO Contributor: Prof Maccon Keane	Page 2 of 3
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		

## REFERENCES:

1. Early Breast Cancer Trialists' Collaborative Group. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. *Lancet*. 2005;365(9472):1687-1717.
2. Davies C, Godwin J, Gray R, et al. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. *Lancet* 2011;378(9793):771-784.
3. Davies C, Pan H, Godwin J, et al. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. *Lancet* 2013;381(9869):805-16.
4. Delozier T, Switsers O, Genot JY et al. Delayed adjuvant tamoxifen: ten-year results of a collaborative randomized controlled trial in early breast cancer (TAM-02 trial). *Ann Oncol* 2000;11:515-9.
5. Eisen A, Trudeau M, Shelley W, et al. Aromatase inhibitors in adjuvant therapy for hormone receptor positive breast cancer: a systematic review. *Cancer Treat Rev* 2008;34(2):157-174.
6. Muss HB, Case LD, Atkins JN, et al. Tamoxifen versus high-dose oral medroxyprogesterone acetate as initial endocrine therapy for patients with metastatic breast cancer: a Piedmont Oncology Association study. *J Clin Oncol* 1994;12(8):1630-1638.
7. Novaldex D® Summary of Product Characteristics. Accessed Oct 2020. Last updated Oct 2020. Available at: [https://www.hpra.ie/img/uploaded/swedocuments/Licence\\_PA1019-014-001\\_08062021092220.pdf](https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA1019-014-001_08062021092220.pdf)

Version	Date	Amendment	Approved By
1	1/11/2014		Prof Maccon Keane
2	20/10/2016	Reviewed no changes	Prof Maccon Keane
3	26/11/2018	Updated with new NCCP regimen template and clarified treatment duration	Prof Maccon Keane
4	10/11/2020	Reviewed	Prof Maccon Keane

Comments and feedback welcome at [oncologydrugs@cancercontrol.ie](mailto:oncologydrugs@cancercontrol.ie).

NCCP Regimen: Tamoxifen Monotherapy	Published: 01/11/2014 Review: 10/11/2025	Version number: 4a
Tumour Group: Breast NCCP Regimen Code: 00253	IHS/ISMO Contributor: Prof Maccon Keane	Page 3 of 3
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician, and is subject to HSE's terms of use available at <a href="http://www.hse.ie/eng/Disclaimer">http://www.hse.ie/eng/Disclaimer</a></p> <p><i>This information is valid only on the day of printing, for any updates please check <a href="http://www.hse.ie/NCCPchemoregimens">www.hse.ie/NCCPchemoregimens</a></i></p>		