

Etoposide and CISplatin 20mg/m² (EP) 5 Day Therapy

INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Treatment of good prognosis (IGCCCG criteria) metastatic germ cell tumours (both non-seminoma and seminoma)	C62	00301a	Hospital

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patient's individual clinical circumstances.

Treatment with etoposide and CISplatin is administered on 5 consecutive days (days 1-5), of a 21 day cycle and repeated for 4 cycles or until disease progression or unacceptable toxicity develops.

Facilities to treat anaphylaxis MUST be present when the chemotherapy is administered.

Admin Order	Day	Drug	Dose	Route	Diluent & Rate
1	1-5	Etoposide	100mg/m ²	IV infusion	1000ml 0.9% NaCl over 60 minutes ^b
2	1-5	CISplatin	20mg/m ²	IV infusion	1000ml 0.9% NaCl over 1 hour (Pre hydration therapy required) ^a
<p>^aPrehydration therapy required for CISplatin See local hospital policy recommendations. Suggested <u>prehydration</u> for CISplatin therapy: 1. Administer 10mmol magnesium sulphate (MgSO₄) ((+/-KCl 20mmol/L if indicated) in 1000 mL sodium chloride 0.9% over 60 minutes. Administer CISplatin as described above</p> <p>^bHypotension following rapid IV administration has been reported. Longer infusion times may be required based on the patient's tolerance</p>					

ELIGIBILITY:

- Indications as above
- ECOG status 0-3

EXCLUSIONS:

- Hypersensitivity to etoposide, CISplatin or any of the excipients.
- CISplatin
 - Pre existing neuropathies ≥ grade 2
 - Creatinine clearance <40mL/min
 - Significant hearing impairment/tinnitus
- Severe liver impairment (etoposide)

NCCP Regimen: EP Therapy	Published: 08/04/2016 Review: 28/11/2024	Version number: 4
Tumour Group: Genitourinary/Gynaecology NCCP Regimen Code: 00301	ISMO Contributor: Prof Maccon Keane	Page 1 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</p>		

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Medical Oncologist.

TESTS:

Baseline tests:

- FBC, renal, liver, creatinine
- Consider sperm banking for appropriate patients prior to initiation of therapy

Regular tests:

- FBC weekly during treatment
- Renal, liver, creatinine prior to each treatment cycle

Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant

Haematological:

- Delay and dose reductions are not recommended as the efficacy of this treatment may be greatly compromised.
- All delays to treatment must be approved by prescribing consultant.
- Prophylactic use of G-CSF is not recommended.
- G-CSF is indicated in patients receiving their second or subsequent cycle of EP who have had an episode of neutropenic fever or who have not recovered their neutrophil count by Day 5.

NCCP Regimen: EP Therapy	Published: 08/04/2016 Review: 28/11/2024	Version number: 4
Tumour Group: Genitourinary/Gynaecology NCCP Regimen Code: 00301	ISMO Contributor: Prof Maccon Keane	Page 2 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Renal and Hepatic Impairment:

Table 2: Dose modification in renal and hepatic impairment

Drug	Renal impairment		Hepatic Impairment			
Etoposide	CrCl (ml/min)	Dose	Bilirubin (micromol/L)		AST (Units/L)	Dose
	>50	100%				
	15-50	75%	>51	or	>180	Clinical decision
	<15	50%				
Subsequent dosing should be based on patient tolerance and clinical effect.						
CISplatin	CrCl (ml/min)	Dose	No dose reduction necessary			
	≥ 60	100%				
	*45-59	75%				
	<45	Hold CISplatin or delay with additional IV fluids				
*Due to the curative intent of this chemotherapy regimen, in cases where CrCl falls between 45-59ml/min it may be appropriate to maintain dose of CISplatin but with extra hydration, longer infusion time and daily Creatinine measurements at the discretion of the prescribing consultant.						

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL:

CISplatin High
Etoposide Low (Refer to local policy).

PREMEDICATIONS:

Hydration prior to CISplatin administration (Reference local policy or see recommendations above).

OTHER SUPPORTIVE CARE:

No specific recommendations

ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

- **Hypersensitivity:** Hypersensitivity reactions have been reported with etoposide and CISplatin. Monitor infusion of etoposide for the first 15 minutes for signs of hypotension.
- **Neutropenia:** Fever or other evidence of infection must be assessed promptly and treated appropriately. Avoid aminoglycoside antibiotics.
- **Renal Toxicity:** Nephrotoxicity is common with CISplatin. Strongly encourage oral hydration. If oral hydration is not possible (e.g. excessive nausea), IV hydration is indicated. Avoid nephrotoxic drugs such as aminoglycoside antibiotics where possible. Where treatment with nephrotoxic drugs must be used, monitor renal function.
- **Ototoxicity and sensory neural damage:** These are associated with CISplatin therapy. They should be assessed by history prior to each cycle.

NCCP Regimen: EP Therapy	Published: 08/04/2016 Review: 28/11/2024	Version number: 4
Tumour Group: Genitourinary/Gynaecology NCCP Regimen Code: 00301	ISMO Contributor: Prof Maccon Keane	Page 3 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</p>		

DRUG INTERACTIONS:

- CISplatin may potentiate the nephrotoxic and ototoxic effects of loop diuretics and aminoglycosides so concurrent use should be avoided.
- Concomitant CISplatin therapy is associated with reduced total body clearance of etoposide.
- CYP3A4 inducers may increase the clearance of etoposide.
- CYP3A4 and p-gp inhibitors may decrease the clearance of etoposide
- Current drug interaction databases should be consulted for more information

ATC CODE:

CISplatin L01XA01
Etoposide L01CB01

REFERENCES:

1. Culine, S., P. Kerbrat, A. Kramar, et al. Refining the optimal chemotherapy regimen for good-risk metastatic nonseminomatous germ-cell tumors: a randomized trial of the Genito-Urinary Group of the French Federation of Cancer Centers (GETUG T93BP). *Ann Oncol* 2007;18(5):917-924.
2. Einhorn LH, Williams SD, Loehrer PJ, et al. Evaluation of optimal duration of chemotherapy in favorable-prognosis disseminated germ cell tumors: a Southeastern Cancer Study Group protocol. *J Clin Oncol* 1989; 7:387-91
3. Cisplatin (Eloxatin®) Summary of Product Characteristics HPRA. Last updated: 11/03/2019. Accessed November 2019 Available at: <https://www.hpra.ie/img/uploaded/swedocuments/Final%20approved%20SPC%20PA0822.199.001.pdf>
4. Etoposide 20 mg/ml Concentrate for Solution for Infusion Summary of Product Characteristics. HPRA Last updated: 29/07/2019 Accessed November 2019 Available at: https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA2059-036-001_29072019103821.pdf
5. Dosage Adjustment for Cytotoxics in Renal Impairment January 2009; North London Cancer Network. Available at <http://londoncancer.org/media/65600/renal-impairment-dosage-adjustment-for-cytotoxics.pdf>
6. Dosage Adjustment for Cytotoxics in Hepatic Impairment January 2009; North London Cancer Network. Available at <http://londoncancer.org/media/65594/hepatic-impairment-dosage-adjustment-for-cytotoxics.pdf>
7. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V1 2018. Available at: <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp%20antiemetic%20classification%20document%20v1%202018.pdf>

NCCP Regimen: EP Therapy	Published: 08/04/2016 Review: 28/11/2024	Version number: 4
Tumour Group: Genitourinary/Gynaecology NCCP Regimen Code: 00301	ISMO Contributor: Prof Maccon Keane	Page 4 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

Version	Date	Amendment	Approved By
1	08/04/2016		Dr Maccon Keane
2	20/09/2017	Updated with new NCCP regimen template	Prof Maccon Keane
3	06/12/2017	Updated with revised Cisplatin hydration regimen recommendations	Prof Maccon Keane
4	20/11/2019	Reviewed. Standardised treatment table and renal dose modifications.	Prof Maccon Keane

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

NCCP Regimen: EP Therapy	Published: 08/04/2016 Review: 28/11/2024	Version number: 4
Tumour Group: Genitourinary/Gynaecology NCCP Regimen Code: 00301	ISMO Contributor: Prof Maccon Keane	Page 5 of 5
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		